


CONFIDENTIAL

iGEM 2010 Progress Report

June 21, 2010






TEAM VECTOR

Aim: biobrick agrobacterium vector

We are working with 6 different plasmids known as pORE:





To achieve our aim we need to remove the MCS and replace it with the biobrick cloning sites

TEAM VECTOR

Progress so far: Backbones

Used restriction digests to remove MCS + promoters/ reporters from each plasmid, leaving the backbone.

- Open Series: cut with *Sac*II and *Spe*I
- Reporter Series: cut with *Hind*III and *Spe*I
- Expression Series: cut with *Hind*III and *Spe*I




TEAM VECTOR

Progress: Inserts for reporter and expression series

Created inserts from primers via PCR on the original agrobacterium vector designed to retain any reporter or promoter already in the vector and add the Biobrick Multiple Cloning Site

- Reporter Series:

- forward: HindIII cut site, Biobrick MCS, first part of appropriate reporter
- reverse: last part of reporter, Nhe cut site



E3 E4 R1 R3

- Expression Series:

- forward: HindIII cut site, first part of promoter
- reverse: last part of promoter, Biobrick MCS, Nhe cut site

TEAM VECTOR

Progress so far: Ligation of inserts and backbones

- Ligated inserts and backbones of expression and reporter series vectors to create biobrick agrobacterium plasmids.
- Transformed into NEB Turbo Competent E. coli cells.
- Picked colonies and made cultures.

TEAM VECTOR

Next steps:

- Create inserts for O1 and O2 by annealing oligos and digest with SacII and NheI.
- Ligate O1 and O2 backbones and inserts.
- Transform resulting O1 and O2 plasmids into E. coli.
- Transform each plasmid into agrobacteria.
- Sequence constructs to check for accuracy

Next: Team Flavor

TEAM FLAVOR

Our Plan:

- Express different flavors in plants
 - Targeting and altering specific metabolic pathways
 - Producing proteins
- Make BioBrick Parts


Flavors:

- Miraculin
 - Turns 'Sour' to 'Sweet'
 - Ordered from Mr. Gene
 - Arrival Pending... (should be early this week)
- Brazzein
 - 500x to 2000x sweeter than sugar (Wikipedia)
 - Ordered from Mr. Gene
 - Arrival Pending... (should be early this week)
- Valencene
 - Orange/Citrus smell, derived from Valencia oranges
 - Extracted from oranges ourselves
- Wintergreen Scent
 - From the 2006 MIT iGEM team
 - Full pathway available in 2010 iGEM kit
- Banana Scent
 - From the 2006 MIT iGEM team
 - Parts of pathway available in 2010 kit

To Do:

Make BioBrick parts out of all enzymes/proteins in the flavour pathways.

Make BioBrick parts out of pORE vector components



Insert the newly created flavour BioBricks into the optimized Arabidopsis vector (courtesy of Team Vector)

Express in plants!



What we have done this week:

1. BioBrick parts from the registry were transformed and expressed in E. Coli:
 - Full and partial wintergreen pathway
 - Very few colonies
 - Parts of banana pathway
 - Expressed components of banana pathway did not grow!
 - PCR of restriction digest yielded products of incorrect size
2. Primers were designed, and PCR performed on components of the e3 pORE vector
 - Promoter
 - Terminator sequence
 - Terminator sequence + Stop codon
 - We are in the process of inserting these components into standard BioBrick vectors (V0120). First attempt at ligating into vector did not work. Result of second ligation attempt will be seen on Monday.


Digestion of PCR Products

1. Ladder
2. Promoter pENTCUP2
3. NOST
4. empty
5. NOST + STOP
6. Plasmid

Ladder 1 2 3 4 5 6




1 2 3 4 5 6



What we have done this week (cont'd):

3. Valencia orange RNA extraction and cDNA synthesis
 - extracted RNA from flavedo of Valencia orange
 - need to remove pstI sites from valencene gene without altering functionality
 - site directed mutagenesis
 - primers have been ordered




Primers for site directed mutagenesis:

Valencene synthase t30a (amino acid 440 Ile -> Ile)
5' gagaaacatttcgtcctacagcagattccatcctagtt 3'

Valencene synthase c1320a (amino acid 10 Thr -> Thr)
5' aggttgtaaaagcagcatcagttatatgcagactcatgga 3'

Team Allergy



Team Allergy Goals

Short Term

Design hp RNA to target GFP ; introduce to Arabidopsis; observe GFP knockdown through qrt pcr/ fluorescence

Design hp RNA to target Fra a1/ LTP1; introduce to strawberry/Arabidopsis; observe knockdown through assays

Long Term (allergen knockdown)

Knockdown of GFP in GFP Arabidopsis through hp RNA interference

Knockdown of Fra a1 (strawberry allergen) & LTP (lipid transfer protein) in Arabidopsis

What We Did...

Monday and Tuesday :

RNA extraction on strawberry
fruit obtained from Shaws
Obtained sufficient mRNA to
yield the following amounts of
cDNA



Sample #	cDNA concentration (ng/uL)	mRNA concentration (ng/uL)	Part of the strawberry
1	441	11	Outer Receptacle
2	1477	36	Receptacle
3	2749	6.5	Inner Receptacle

PCR

Wednesday and Thursday:

Tried to PCR out Fraa 1 for biobrick cloning
using cDNA from strawberries


Goal: 300bp DNA sequence (Fraa1)

Result?

Unsuccessful --- Primer Dimers? PCR products were ~ 100 bp

What went wrong? We eliminated the following possibilities through further testing

- Incorrect annealing temperature by using a temperature gradient
- Too much cDNA in the reaction by reducing the amount of cDNA from micrograms to ~100 nanograms/ 50 uL reaction



One Important Question!

Was there mRNA coding for Ffraa1 in the strawberries when we extracted mRNA?

Further research on Ffraa1 reveals that Ffraa1 is part of the ripening mechanism of strawberries

Strawberries are non-climactic fruit, which stop all ripening metabolic processes after removal from plant. Our strawberries were store-bought, and therefore have been “dead” for at least two or three days, giving the mRNA coding for Ffraa1 plenty of time to degrade.

Next experiment ----

- Make sure that we have reason to believe that our sample is in a state that contains the appropriate mRNA

Onward!

Search for allergens in arabidopsis



It should be ready(er) by now.

Namely:

LTP1 (documented allergen)

Bet v 1 (thaliana <= latera <= birch homologue via BLAST)

Others (list, links on wiki)






*Bet v I: Pathogenesis-related proteins

*Lipid transfer protein



*Germin: stress protein

*Profilin: actin binding





081729
9 vi 10
10 SE

081733
9 vi 10

081733
9 vi 10
12

081723
9 vi 10
12

081733
9 vi 10
11

081733
9 vi 10
11

584733
9 vi 10

584733
9 vi 10
11

ca 84733
9 vi 10

11

ca 84733
9 vi 10
11



50ug isn't much, but
neither are our plants!



We would feel much less
guilty sampling plants of
this size or larger.

More importantly,
sampling should be more
reliable.




They should be larger by now.

After all, they spent the weekend happily growing in the underground greenhouse bunker.







Future Directions




TEAM GENETIC FENCE





Methoxyfenozide



DEATH







Last Week

- Methoxyfenozide was ordered and arrived Thursday
- Made requests to other labs for LacI with NLS, ActIn promoter with lacO sites (A2pot), and Cassava Vein Mosaic Virus promoter, as well as plasmids containing the entire monopartate and bipartate switch mechanisms
- Barstar and Barnase ordered from ADDGENE


Last Week, Cont.

- VP16, GAL4 DNA Binding Domain, and LacI in biobricks registry
- LacI
 - Two biobricks in registry, one wildtype and one including and LVA rapid degradation tail
 - Transformed, minipreped and digested both LacI and LacI+LVA
- VP16
 - Transformed, minipreped and digested VP16
- GAL4 DBD
 - Transformed
- One run of PCR to attach NLS to LacI
 - Also annealed oligos to form NLS alone for biobrick


Alternate/Secondary System: Crelox and IPTG

- Attractive paradigm (lactose analog instead of pesticide)
- However untested and relies on less-characterized parts

Lac +
Cre/Lox



Root +
LacI



External control of transgene expression in tobacco plastids using the bacterial *lac* repressor

Stefan K. Mühlbauer and Hans-Ulrich Koop*

Icon Genetics AG, Research Centre Freising, Lise-Meitner-Str. 30, 85354 Freising, Germany

Received 17 March 2005; revised 9 June 2005; accepted 17 June 2005.

*For correspondence (fax +49 89 17861165; e-mail koop@lrz.uni-muenchen.de).

Summary


Although several induction systems have been described for plants containing transgenes in the nucleus, to date there is only one method for controlling transgene expression in plastids. This consists of chemical induction of a nuclear gene and import of the gene product into plastids, so that transformation of two cellular compartments is required. Here we describe a system for external control of plastid gene expression which is based entirely on plastid components and can therefore be established in a single transformation step. Our system uses modified promoters containing binding sites for the bacterial *lac* repressor. Chemical induction can be made with intact plants or after harvesting, which provides ecological and economic benefits.

Keywords: plastid transformation, transgene expression, induction, *lac* repressor, IPTG.

(b) - + IPTG



- + IPTG



Control of plastid gene expression 943

(a)

PrrnP1: ..tggataagaggctcgtggga -35 ttgacgtgagggggcagggatggc -10 tatatttctgggagcgaactccgggcg..


Prrnlac1: ..tggataagaggctcgtggga -35 ttgacgtgagggggcagggatggc -10 tatatttctggga attgtgagcggataacaag..

Prrnlac2: ..tggataagaggctcgtggga -35 ttgacg attgtgagcggataaca -10 tatatttctgggagcgaactccgggcg..

Prrnlac3: ..tggataagaggctcgtggga -35 ttgacg attgtgagcggataaca -10 tatatttctggga attgtgagcggataacaag..

Plac: ..actcattaggcaccacaggc -35 ttta -10 ctttatgcttcgggctcgtatgttgtgtgga attgtgagcggataacaat..

Tissue Specific Promoters



Back up:

Arabis thaliana	All	Target	Base
callus	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
cell culture / primary cell	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
sperm cell	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
seedling	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
cotyledons	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
hypocotyl	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
radicle	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
imbibed seed	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
inflorescence	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
flower	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
carpel	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
ovary	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
stigma	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
petal	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
sepal	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
stamen	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
pollen	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
abscission zone	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
pedicel	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	
silique	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
seed	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
embryo	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
endosperm	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
micropylar endosperm	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
peripheral endosperm	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
chalazal endosperm	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
testa (seed coat)	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
general seed coat	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
chalazal seed coat	<input type="checkbox"/>	<input checked="" type="checkbox"/>	
suspensor	<input type="checkbox"/>	<input checked="" type="checkbox"/>	

Other considerations

- Plastid DNA and Chloroplast modification
 - dramatically reduces risk of cross-pollination
 - Increases technical difficulty of transformation (agrobacterial vector no longer sufficient)
- Time specific tissue-specific promoters for cre-lox?